DO NOT OPEN THIS TEST BOOKLET UNTIL YOU ARE ASKED TO DO SO

TEST BOOKLET SERIES

TEST BOOKLET PGT(MATHEMATICS)-2016



Time Allowed : 2 Hours		[Maximum	100
	All questions carry equal marks		

INSTRUCTIONS

- Immediately after the commencement of the examination, you should check that test booklet does not have any unprinted or torn or missing pages or items, etc. If so, get it replaced by a complete test booklet.
- 2. Encode clearly the test booklet series A, B, C or D as the case may be in the appropriate place in the answer-sheet.
- 3. Write your Roll Number only in the box provided alongside.

 Do not write anything else on the Test Booklet.
- This Test Booklet contains 100 items (questions). Each item comprises four responses (answers). Choose only one response for each item which you consider the best.
- 5. After the candidate has read each item in the Test Booklet and decided which of the given responses is correct or the lest, he has to mark the circle containing the letter of the selected response by blackening it completely with Black or Blue ball pen. In the following example, response "C" is so marked:
 - (A) (B) (D)
- 6. Do the encoding carefully as given in the illustrations. While encoding your particulars or marking the answers on answer sheet, you should blacken the circle corresponding to the choice in full and no part of the circle should be left unfilled. After the response has been marked in the ANSWER SHEET, no erasing/fluid is allowed.
- You have to mark all your responses ONLY on the ANSWER SHEET separately given according to 'INSTRUCTIONS FOR CANDIDATES' already supplied to you. Responses marked on the Test Booklet or in any paper other than the answer sheet shall not be examined.
- All items carry equal marks. Attempt all items. Your total marks will depend only on the number of correct responses marked by you in the Answer Sheet. There will be no negative marking.
- Before you proceed to mark responses in the Answer Sheet fill in the particulars in the front portion of the Answer Sheet as per the instructions sent to you.
- If a candidate gives more than one answer, it will be treated as a wrong answer even if one of the given answers happens to be correct.
- 11. After you have completed the test, hand over the Answer Sheet only, to the Invigilator.

PGT(MATHEMATICS)-2016

Time Allowed: 2 Hours

- Maximum Marks: 100
- If $u = \sin^{-1} \frac{x + 2y + 3z}{x^8 + y^8 + z^8}$, then the value of $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z}$ is: 1.
 - (A) tan u

(B) 5 tan u

(C) 6 tan u

- (D) −7 tan u
- If u = f(2x 3y, 3y 4z, 4z 2x), then the value of $\frac{1}{2} \frac{\partial u}{\partial x} + \frac{1}{3} \frac{\partial u}{\partial y} + \frac{1}{4} \frac{\partial u}{\partial z}$ 2.
 - $(A) \quad x + y + z$

is :

(B) 2x - 3y + z

- (D) 0
- The radius of curvature at the point (3a/2, 3a/2) of the curve $x^3 + y^3 = 3axy$ 3. is:
 - (A) $\frac{3a}{\sqrt{2}}$. (B) $\frac{3a}{2\sqrt{2}}$

- (D) $\frac{3a}{8\sqrt{2}}$
- The value of the integral $\int_0^{2a} x^3 \sqrt{2ax x^2} dx$ is : 4.
 - (A) $\frac{7\pi a^5}{8}$

(B) $\frac{7\pi a^5}{9}$

(C) $\frac{7\pi a^5}{10}$

(D) $\frac{7\pi a^5}{11}$

5.	The area of the cardioid $r =$	$= a(1 - \cos \theta)$ is:	
	(A) $\frac{3\pi a^2}{2}$	(B) $\frac{\pi a^2}{3}$	٠
	(C) $\frac{3\pi a^2}{4}$	(D) 3πa ²	
6.	The area of the plane regio	on bounded by the parabola $y^2 = 4ax$ and	d the
	latus rectum, is :		
	(A) $\frac{2}{3}a^2$	(B) $\frac{4}{3}a^2$	
	(C) $\frac{5}{3}a^2$	(D) $\frac{8}{3}\alpha^2$	
7.	The surface of the solid for	med by revolving the cardioid $r = a(1 +$	cos θ)
	about the initial line is:		
	(A) $\frac{\pi a^2}{5}$	(B) $\frac{16\pi a^2}{5}$	
	(C) $\frac{32\pi a^2}{5}$	(D) $\frac{48\pi a^2}{5}$	
8.	The semi-vertical angle of	a cone of maximum volume and of given	slant
	height is:		
	(A) $\tan^{-1}\sqrt{3}$	(B) $\tan^{-1}\sqrt{5}$	
	(C) ton-1 /2	(D) tan ⁻¹ √6	

PGT(MATHEMATICS)-2016—A

P.T.O.

The value of the integral $\int \frac{dx}{x(x^n+1)}$ is:

$$(A) \quad \frac{1}{n} \log \frac{x^n}{x^n + 1} + c$$

(B)
$$\frac{1}{n}\log\frac{x^n+1}{x^n}+c$$

(C)
$$\frac{1}{n}\log x^n + c$$

$$(D) \quad \frac{1}{n} \log \frac{1}{x^n + 1} + c$$

If $I_n = \int_0^{\pi/3} \tan^n x \, dx$, then: 10.

$$(\mathbf{A}) \quad \left(n+1\right) \left(\mathbf{I}_n - \mathbf{I}_{n-2}\right) = \left(\sqrt{3}\right)^{n-1} \qquad (\mathbf{B}) \quad \left(n-1\right) \left(\mathbf{I}_n + \mathbf{I}_{n-2}\right) = \left(\sqrt{3}\right)^{n-1}$$

(B)
$$(n-1)(I_n + I_{n-2}) = (\sqrt{3})^{n-1}$$

(C)
$$(n+1)(I_n + I_{n-2}) = (\sqrt{3})^{n+1}$$
 (D) $n(I_n + I_{n-2}) = \sqrt{3}$

(D)
$$n(I_n + I_{n-2}) = \sqrt{3}$$

The value of the integral $\int_0^{\pi/6} \cos^6 3\theta \sin^2 6\theta d\theta$ is : 11.

$$(A) \quad \frac{\pi}{384}$$

(B)
$$\frac{5\pi}{384}$$

(C)
$$\frac{7\pi}{384}$$

(D)
$$\frac{9\pi}{384}$$

The value of the double integral $\iint xy \, dx \, dy$ over the region in the positive 12. quadrant for which $x + y \le 1$, is :

(A)
$$\frac{1}{6}$$

(B)
$$\frac{1}{12}$$

(C)
$$\frac{1}{18}$$

(D)
$$\frac{1}{24}$$

PGT(MATHEMATICS)-2016-

13.	The equation of the plane through	the point (-1, 2, 4) and parallel to the
	plane $2x + 3y - 5z + 6 = 0$ is:	
	(A) $2x + 3y - 5z + 16 = 0$	(B) $2x + 3y - 5z + 17 = 0$
	(C) $2x + 3y - 5z + 18 = 0$	(D) $2x + 3y - 5z + 19 = 0$

14. The coordinate of the point where the line joining the points (2, -3, 1) and (3, -4, -5) cuts the plane 2x + y + z = 7 is:

(A) (1, -2, 7)

(B) (1, 2, 7)

(C) (-1, 2, -7)

(D) (-1, -2, 7)

15. The volume of tetrahedron whose vertices are (a, 1, 2), (3, 0, 1), (4, 3, 6) and (2, 3, 2) is:

(A) 6 - a

(B) 6 - 2a

(C) 6 - 3a

(D) 6 - 4a

16. The centre and radius of the sphere $x^2 + y^2 + z^2 - 2x + 4y - 6z = 11$ are:

(A) (1, -2, 3) and 5

(B) (1, -2, -3) and 5

(C) (1, -2, -3) and 15

(D) (-1, -2, -3) and 25

17. The equation of the tangent plane to the central conicoid $3x^2 - 5y^2 + z^2 + 2 = 0 \text{ at the point } (1, 1, 0) \text{ is }:$

(A)
$$2x + 5y - 7 = 0$$

(B)
$$3x - 5y + 2 = 0$$

(C)
$$x + 5y - 6 = 0$$

(D)
$$x + y - 2 = 0$$

18. The angle between the surfaces $x \log z = y^2 - 1$ and $x^2y = 2 - z$ at the point (1, 1, 1) is:

(A)
$$\cos^{-1} \frac{1}{\sqrt{30}}$$

(B)
$$\sin^{-1}\frac{1}{\sqrt{30}}$$

(C)
$$\tan^{-1} \frac{1}{\sqrt{30}}$$

(D)
$$\cot^{-1} \frac{1}{\sqrt{30}}$$

19. The inverse Laplace transform of the function $\frac{s}{(s^2 + a^2)^2}$ is :

(A)
$$\frac{t \sin at}{2a}$$
.

(B)
$$\frac{t \cos at}{2a}$$

(C)
$$\frac{\sin at + \cos at}{2a}$$

(D) sin at cos at

20. The Fourier cosine transform of $\frac{1}{1+x^2}$ is :

(A)
$$\frac{\pi}{2}e^{-s}$$

PGT(MATHEMATICS)-2016-A

- 21. The inverse z-transform of the function $\frac{z}{(z+2)(z+3)}$ is:
 - (A) $(-2)^n + (-3^n)$

(B) $(-2)^n - (-3^n)$

(C) $(-2)^n - 3^n$

- (D) 2ⁿ 3ⁿ
- 22. The value of the $\lim_{x\to 0} \frac{(1+x)^{1/x}-e}{x}$ is:
 - (A) $\frac{e}{3}$

(B) $-\frac{e}{2}$

(C) $\frac{e}{4}$

- (D) e
- 23. The function $f(x) = \sin x + \cos x$ is:
 - (A) increasing in interval (0, 5π/4)
 - (B) decreasing in interval $(0, \pi/4)$
 - (C) decreasing in interval $(5\pi/4, 2\pi)$
 - (D) decreasing in interval (π/4, 5π/4)

24. The maximum and minimum value of the function $f(x) = \sin x + \frac{1}{2} \sin 2x + \frac{1}{2} \sin 2x$

$$\frac{1}{3}\sin 3x$$
 for all $x \in [0, \pi]$ are:

(A) $\frac{4\sqrt{2}+3}{6}$ and $\frac{\sqrt{3}}{4}$ respectively

(B) $\frac{4\sqrt{2}-3}{6}$ and $\frac{\sqrt{3}}{4}$ respectively

(C) $\frac{4\sqrt{2}-3}{6}$ and $\frac{\sqrt{5}}{4}$ respectively

(D) $\frac{4\sqrt{2}-3}{2}$ and $\frac{\sqrt{3}}{2}$ respectively

25. In a plane triangle ABC, the maximum value of cos A cos B cos C is :

(A) 1/2

(B). 1/4

(C) 1/6

D) 1/8

26. The volume of the greatest rectangular parallelopiped that can be inscribed

in the ellipsoid $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$ is :

(A) 8abc

(B) 8abc/3

(C) 8abc/√3

(D) 8abc/(3\sqrt{3})

PGT(MATHEMATICS)-2016-A

- 27. The function $f(x) = x^2$ is:
 - (A) uniformly continuous on [0, ∞)
 - (B) uniformly continuous on (-∞, 0]
 - (C) uniformly continuous on (-∞, ∞)
 - (D) uniformly continuous on every closed and finite interval
- 28. The function $f(x) = \frac{2[x]}{3x |x|}$
 - (A) is continuous at x = 1 and discontinuous at x = -1/2
 - (B) is discontinuous at x = 1 and continuous at x = -1/2
 - (C) is continuous at x = -1/2, The pexams.in
 - (D) is discontinuous at x = -1/2, 1
- 29. The function $f(x) = \frac{x |x|}{x}$ is not continuous at:
 - $(A) \quad x = 0$

(B) x = 1

(C) x = 2

- (D) x = 3
- 30. Rolle's theorem does not holds for the function :
 - (A) $f(x) = 1 (x 1)^{2/3}$ on [0, 2]
 - (B) $f(x) = \sqrt{1-x^2}$ on [-1, 1]
 - (C) $f(x) = x^3 4x$ on [-2, 2]
 - (D) $f(x) = \cos x$ on $[-\pi/2, \pi/2]$

31.	For	the set $G = \{-2, -3/2, -4/3, \dots \}$, which one of the statements is true ?
	(A)	G is unbounded
	(B)	inf G = -2 and sup G = ∞
	(C)	inf $G = -3/2$ and sup $G = -1$
	(D)	inf $G = -2$ and $\sup G = -1$
32.	Whi	ch one of the following inequalities is not true?
**	(A)	$(1 + x) < e^x < 1 + xe^x$ for all $x > 0$
	(B)	$\frac{x}{1+x} < \log(1+x) < x \text{ for all } x > 0$
	(C)	$ \tan^{-1} x - \tan^{-1} y \le x - y \text{ for all } x, y \in \mathbf{R}$
	(D)	$ \sin x - \sin y \ge x - y $ for all $x, y \in \mathbf{R}$
33.	Whi	ich one of the following set is not closed?
	(A)	R
	(B)	Q
	(C)	empty set
	(D)	every closed and finite interval
34.	Wh	ich one of the following is countable ?
	(A)	$\mathbf{R} \setminus \mathbf{Q}$ (B) $\mathbf{Q} \setminus \mathbf{Z}$
	(C)	$\mathbf{R} \setminus \mathbf{Z}$ (D) $\mathbf{R} \setminus \mathbf{N}$
PGT	(MAT	HEMATICS)-2016—A 10

Which one of the set is open in R ? (B) R\Z (A), Z $R \setminus Q$ (C) Q Which one of the following statements is not true ? 36. Every point of the set Q of rational is a limit point Every point of the closed interval [a, b] is its limit point (C) a finite set has at least one point (D) $\left\{\frac{1}{n}: n \in \mathbb{N}\right\}$ has only one limit point The sequence $\left(1+\left(-1\right)^n\right)$: 37. diverges (A) converges oscillate infinitely (C) oscillate finitely An infinite series $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{1}{n}$ converges to log 3 converges to log 2

39. The sum of an infinite series $\frac{5}{1.2.3} + \frac{7}{3.4.5} + \frac{9}{5.6.7} + \dots$ is:

(A) 3 log 2 - 1

(B) log 2 - 1

(C) log 2

(D) 3 log 2

40. The series $\sum \left[\frac{1}{n} + \frac{(-1)^n}{\sqrt{n}}\right]$ is :

(A) convergent

(B) divergent

(C) oscillate finitely

(D) oscillate infinitely

41. The solution of the differential equation $\frac{dy}{dx} = \frac{\sin y + y \sin x}{\cos x - x \cos y}$ is :

(A) $y \cos x + x \sin y = c$

(B) $2y \cos x - x \sin y = c$

(C) $y \cos x - x \sin y = c$

(D) $x \cos x - y \sin x = c$

42. The general solution of the differential equation $(D-2)^3y = e^{2x}$ is :

(A)
$$(A + Bx + Cx^2 + x^2/4)e^{2x}$$

(B)
$$(A + Bx + Cx^2 + x^2/6)e^{2x}$$

(C)
$$(A + Bx + Cx^2 + 1/4)e^{2x}$$

(D) $(A + Bx + Cx^2 + x/4)e^{2x}$ PGT(MATHEMATICS)-2016—A 12 The particular solution of the differential equation $(D^2 - 1)y = x^2 \cos x$ is :

$$(A) \quad x \sin x + (1 - x^2) \cos x$$

(B)
$$x \sin x + (1 - x^2) \frac{\cos x}{3}$$

(C)
$$x \sin x + (1 - x^2) \frac{\cos x}{2}$$
 (D) $x \sin x + (1 - x^2) \frac{\cos x}{4}$

(D)
$$x \sin x + (1 - x^2) \frac{\cos x}{4}$$

Which one of the following is an elliptic partial differential equation ? 44.

(A)
$$\frac{\partial^2 z}{\partial x^2} - 6 \frac{\partial^2 z}{\partial x \partial y} + \frac{\partial^2 z}{\partial y^2} = 0$$

(B)
$$\frac{\partial^2 z}{\partial x^2} + 2 \frac{\partial^2 z}{\partial x \partial y} + \frac{\partial^2 z}{\partial y^2} = 0$$

(C)
$$\frac{\partial^2 z}{\partial x^2} - \frac{\partial^2 z}{\partial x \partial y} + \frac{\partial^2 z}{\partial y^2} = 0$$

(D)
$$\frac{\partial^2 z}{\partial x^2} - 4 \frac{\partial^2 z}{\partial x \partial y} + \frac{\partial^2 z}{\partial y^2} = 0$$

The general solution of the partial differential equation (y + z)p + (z + x)q45. = x + y is:

(A)
$$\phi\left(\frac{x-y}{x-z},(x-y)^2(x-y-z)\right)=0$$

(B)
$$\phi\left(\frac{x-y}{x-z},(x-y)^2(x-y+z)\right)=0$$

(C)
$$\phi\left(\frac{x-y}{x-z},(x-y)^2(x+y+z)\right)=0$$

(D)
$$\phi\left(\frac{x-y}{x+z},(x-y)^2(x+y-z)\right)=0$$

46. The cube roots of the complex number -i are :

(A)
$$i$$
, $\pm \frac{\sqrt{3}}{2} - \frac{1}{2}i$

(B)
$$-i, \pm \frac{\sqrt{3}}{2} - \frac{1}{2}i$$

(C)
$$i, \pm \frac{\sqrt{3}}{2} + \frac{1}{2}i$$

(D)
$$i, 1, \frac{\sqrt{3}}{2} + \frac{1}{2}i$$

47. The function $f(z) = e^z$; $z \in \mathbb{C}$ is:

(A) bounded

(B) increasing

(C) periodic

(D) decreasing

48. Which of the following complex valued function is not analytic?

(A) cos z

 $(B) \sin z$

(C) e²

(D) tan z

49. The complex valued function log z has a branch point at :

 $(A) \quad z = 0$

(B) z = 1

(C) z = 2

(D) $z = \infty$

50. The value of the contour integral $\oint_{|z|=1} \frac{dz}{2-\overline{z}}$ is :

(A) πi

(B) 2πi

(C) ni/2

(D) 0

PGT(MATHEMATICS)-2016—A

51. The radius of the convergence of the power series $\sum_{n=1}^{\infty} \frac{(-1)^{n+1} n}{5n^2 + 3} (z - z_0)^n$

is ;

(A) 0

(B) 1

(C) 2

(D) 3

52. For the complex valued function $f(z) = \frac{e^z}{z - \sin z}$, the point z = 0 is :

(A) a branch point

(B) removable singularity

(C) an essential singularity

(D) a pole of order 3

53. The bilinear mapping which maps the upper half of the z-plane on to the right half of the w-plane is:

(A)
$$w = \frac{2iz}{z+1}$$

$$(B) \quad w = \frac{2z}{z-1}$$

(C)
$$w = \frac{2i}{z+1}$$

(D)
$$w = \frac{iz}{2z+1}$$
.


54. The bilinear transformation $w = \frac{az+b}{cz+d}$; $ad-bc \neq 0$ is not conformal at:

$$(A) z = c$$

$$(B) z = d$$

(C)
$$z = 0$$

 Under the mapping w = 1/z, the image of the circle z - 3 = 5 is a : (A) hyperbola (B) circle (C) square (D) strip Let A(1, -2), B(-3, 4) and C(2, 2) be the three vertices of the triangle ABC. Then the length of the median from C to the side AB is : (A) √5 (B) √10 (C) √15 (D) √20 The trace of a 3 matrix is 2. Two of its eigen values are 1 and 2. Then the third eigen value is : (A) -1 (B) 0 (C) 1 (D) 2 The matrix									
	(A) hyperbo	la			(B)	circle			
	(C) square				(D)	strip			
56.	Let A(1, -2),	B(-3, 4) ar	nd C(2	2, 2) be	the t	three vertic	es of the t	riangle AI	3C
	Then the len	gth of the	medi	an from	n C t	to the side	AB is:		
	(A) √5			*2	(B)	$\sqrt{10}$			
	(C) $\sqrt{15}$				(D)	$\sqrt{20}$			
57.	native at the second		is 2.	Two of	its ei	gen values	are 1 and	2. Then t	he
	(A) -1				(B)	0			
	(C) 1				(D)	2			
	,	1 1	3				* 5		
58.	The matrix	5 2	6	is:					
		-2 -1	-3)				90	9	
	(A) an invol	utary			(B)	nilpotent			
	(C) an idem	potent			(D)	skew-symr	netric		
PGT(I	MATHEMATICS)-2016—A		16		*			



62.	The	variance of 20 observations is 5.	If ea	ch observation is multiplied by 2,
	then	the new variance of the resulti	ing o	bservation is:
	(A)	15	(B)	20
	(C)	25	(D)	30
63.	be :	ne of the regression coefficient	is g	reater than 1, then other must
		greater than 2	(B)	less than 1
	(C)	equal to 1	(D)	equal to 2
64.	A di	e is tossed. If the number is odd	, the	n the probability that it is prime,
	(A)	$\frac{1}{4}$	(B)	$\frac{1}{2}$
	(C)	$\frac{2}{3}$	(D)	$\frac{3}{4}$
65.	The	set $S = \{1, 5, 7, 11\}$ is a g	roup	with respect to multiplication
	mod	ulo :		₱.
	(A)	5	(B)	7
	(C)	9	(D)	12
PGT(MATE	HEMATICS)-2016—A 18		

66.	In the multiplicative group $G = \{1, -1, i\}$, -i, the order of an elemen
	<i>−i</i> is :	
Y 8	(A) 1 (B) 2	
	(C) 3 (D) 4	
67.	For a finite group G, Lagrange theorem st	ates that the :
*	(A) order of an element divides order of (3.
	(B) order of any subgroup divides order of	f G
	(C) sum of orders of elements in G is the	order of G
	(D) order of G equals order of identity ele	ement
68.	If H_1 , H_2 are subgroups of a group G , the	n which one of the following is
	a subgroup of G?	
	(A) H ₁ U H ₂ (B) H	$H_1 \cap H_2$
	(C) H ₁ \H ₂ (D) H	I_1H_2
69.	The quotient group is always defined excep	ot possibly when the group is
	(A) abelian (B) c	yelic
		A
	(C) a group of prime order (D) n	on-abelian

	multiplication ?		
	(A) the set of even integers	2.0	
	(B) the set of integers which are	multiple of 3	
	(C) the set of positive integers		
	(D) the set of integers		
71.	In the ring \mathbf{Z}_{10} , the divisor of 0	is:	
	(A) 1	(B) 2	
	(C) 3	(D) 7	
72.	Consider A = $\{q \in \mathbf{Q} : q^2 \le$	2) as a subset of	(\mathbf{Q}, d) , where
	d(x, y) = x - y . Then A is:		
70	(A) closed but not open in Q		
	(B) open but not closed in \mathbf{Q}		
	(C) neither open nor closed in Q	P -	
	(D) both open and closed in \mathbf{Q}		
73.	The set of all integers Z consi	dered as a subspace	of (\mathbf{R}, d) where
	d(x, y) = x - y , is :	* * * * * * * * * * * * * * * * * * *	
	(A) closed but not complete	(B) complete but n	not closed
	(C) both closed and complete	(D) neither closed	nor complete
PGT	T(MATHEMATICS)-2016—A	20	net in

Which one of the following is not a ring with respect to usual addition and

70.

74.	4. The closure Y of a totally bounded subset Y of a metric space X :					
	(A)	is totally bounded				
	(B)	may not be totally	bounded even if	X is complete		
	(C)	is totally bounded if	and only if X	is complete		
	(D)	is totally bounded if	and only if X	is compact	IN THE	
75.	Let	X, Y be metric spaces	and let $f: X \rightarrow$	Y be a continuous for	unction. Then	
	the	image f(A) of a boun	ded subset A of	X is bounded:		
	(A)	always	(B)	if A is also open		
	(C)	if A is compact	(D)	if A is complete	No.	
76.	Let	U be subset of connec	ted metric space	X which is both ope	n and closed.	
	The	n U is:				
	(A)	neither ϕ nor X	(B)	x	4.	
	(C)	•	(D)	either ϕ or X		
77.	The	image of a connecte	d metric space	under a continuous	real-valued	
	func	tion is:			-	
	(A)	R	(B)	a bounded subset o	f R	
	(C)	an interval in R	(D)	not an interval in	R	
PGT(I	MATH	IEMATICS)-2016—A	21	72 ×	P.T.O.	
			112			

- 78. Let X be a normed linear space. Then which one of the following is false?
 - (A) If X has a Schauder basis, then X is separable
 - (B) Every separable space has Schauder basis
 - (C) l₁ has a Schauder basis
 - (D) l2 has a Schauder basis
- 79. Which one of the following is false?
 - (A) A compact subset of a metric space is closed and bounded
 - (B) A closed and bounded subset of metric space is compact
 - (C) A compact subset of a finite dimensional normed space is closed and bounded
 - (D) A closed and bounded subset of a finite dimensional normed space is compact
- 80. The dual of l_p is itself if :

(A)
$$p = 1$$

(B)
$$p = \frac{1}{2}$$

(C)
$$p = 2$$

81.	1. Which lake is the source of Parbati river of Kullu ?						
	(A)	Surajtal	. 4	- P.	(B)	Sketi	
A.	(C)	Mantilai		A	(D)	Karali	
82.	Whi	ch river's tril	outaries ar	e Malaha	t Kh	ad, Garni Khad a	and Hum Khad?
	(A)	Beas		3	(B)	Swan	
÷	(C)	Giri	Ta ((D)	Pabbar	
83.	To w	vhich ancient	tribe did K	ing Sham	bar,	who fought agains	t the Aryan King
	Divo	odas, belong	?			±#	
, 1	(A)	Kinnars	>		(B)	Khasas	
	(C)	Kirats		v V	(D)	Dasas	
84.	Who	is the auth	or of Cata	logue of	the B	thuri Singh Muse	um, Chamba ?
-	(A)	J. Ph. Voge	Ï.		(B)	G.T. Vigne	
	(C)	C.F. Massy		4	(D)	J.B. Lyall	
PGT(MATH	HEMATICS)-20	16—A	23			P.T.O.

85.	The Kullu Princely State under the	ne rajas consisted of seven Waziris. In which
	one of them was Kullu proper i	ncluded ?
	(A) Rupi	(B) Lag Sari
	(C) Parol	(D) Lag Maharaj
86.	Which place near Dharamsala	town of H.P. is called 'Little Israel'?
	(A) Sidhbari	(B) Mataur
	(C) Shahkot	(D) Dharamkot
87.	Which cement company has a	big cement plant at Barmana in Bilaspur
	District of H.P. ?	
	(A) Ambuja	(B) CCI
	(C) Japee	(D) ACC
88.	At which place in Shimla is t	he office of Chief Information Commissioner
	of H.P. ?	
	(A) Richmond Villa	(B) Del Villa
	(C) Revenswood	(D) Armsdale Building
PC	T(MATHEMATICS)-2016—A	24

89.	Around which year was Bishop Co	tton Sch	ool, Shimla shift	ed from its	original
	site, Jutogh ?			2	
	(A) 1846	(B)	1866		
	(C) 1876	(D)	1886	**	P T
90.	In which of the following Distric	cts of H	.P. Kishori Sha	kti Yojan	a is not
	functioning ?				
	(A) Sirmaur	(B)	Kinnaur	9	
	(C) Solan	(D)	Una		
91.	With which sport is OP Jaisha a	ssociate	d ?		
	(A) 800 mtr. race	(B)	1500 mtr. race		
	(C) 20 km walk	(D)	Marathon		
92.	Who is Farooq Khan?				
	(A) Leader of All Party Hurriya	at Confe	rence		,
	(B) Leader of National Conferen	nce			
	(C) Administrator of Lakshadwe	ep Unio	n Territory		
	(D) None of the above				
PGT	(MATHEMATICS)-2016—A	25			P.T.O.

93.	Which day is observed in India as Nirbhaya Day?						
	(A)	July 14	(B)	October 01			
	(C)	November 14	(D)	December 16			
94.	Which state of India is also known as Indrakil, the garden of Indra, the wa						
	god	*					
	(A)	H.P.	(B)	Uttarakhand			
	(C)	Sikkim	(D)	J & K			
95.	Whi	Which railway station in North India is named after a Sufi saint?					
	(A)	Faridabad	(B)	Gaziabad			
	(C)	Ferozabad	(D)	Nizamuddin			
96.	Where is Raqqa, the stronghold of Islamic State ?						
	(A)	A city in Iraq	(B)	A town in Afghanistan			
	i i i i i i i i i i i i i i i i i i i	A city in Iran		A city in Syria			
PGT	MAT	HEMATICS)-2016—A 26					

97.	Whe	en did the People's	Republic of F	Kampuchea change	its name to
911	Cam	nbodia ?		*	
	(A)	1970	(B)	1975	* .
	(C)	1989	(D)	1992	
98.	Who	was crowned Miss	Universe 2015 ?		
	(A)	Pia Alonzo Wurtzba	ach (B)	Aniporn Chalermb	ouranawong
2,	(C)	Ariadna Gutierrez	(D)	Olivia Jordan	\$4.5° 1.5°
99.	Whi	ch Chinese city is th	e venue of G-20	Summit to be held	in September
	2016	5 ?		.100	, K
** .	(A)	Shanghai	(B)	Shenzhen	ş.
•	(C)	Hangzhou	(D)	Biejing	-
100.	At v	which place was Mot	her Teresa born	?	and the second
	(A)	Bitola	(B)	Skopje	
	(C)	Tirana	(D)	Tetovo	
PGT(MATI	HEMATICS)-2016—A	27		P.T.O.